

1. Prove by induction that  $\sum_{k=1}^n \frac{1}{(2k-1)(2k+1)} = \frac{n}{2n+1}$  for  $n = 1, 2, \dots$

Basis:  $n=1$ : L.H.S. =  $\frac{1}{(2-1)(2+1)} = \frac{1}{3}$  R.H.S. =  $\frac{1}{2+1} = \frac{1}{3}$   $\checkmark$

For  $n > 1$  assume  $\sum_{k=1}^{n-1} \frac{1}{(2k-1)(2k+1)} = \frac{n-1}{2(n-1)+1} = \frac{n-1}{2n-1}$

Then  $\sum_{k=1}^n \frac{1}{(2k-1)(2k+1)} = \sum_{k=1}^{n-1} \frac{1}{(2k-1)(2k+1)} + \frac{1}{(2n-1)(2n+1)}$   
 $= \frac{n-1}{2n-1} + \frac{1}{(2n-1)(2n+1)} = \frac{(n-1)(2n+1)+1}{(2n-1)(2n+1)} = \frac{2n^2+n-2n-1+1}{(2n-1)(2n+1)}$   
 $= \frac{2n^2-n}{(2n-1)(2n+1)} = \frac{n \cancel{(2n-1)}}{\cancel{(2n-1)}(2n+1)} = \frac{n}{2n+1}$   $\checkmark$

2. Prove by induction that  $3^n \geq 1 + 2^n$  for  $n = 1, 2, \dots$

Basis:  $n=1$  L.H.S. = 3 R.H.S. =  $1+2=3$   $\checkmark$

For  $n > 1$  assume  $3^{n-1} \geq 1 + 2^{n-1}$

Then  $3^n = 3^{n-1} \cdot 3 \geq (1 + 2^{n-1}) \cdot 3 = 3 + 2^{n-1} \cdot 3 > 1 + 2^{n-1} \cdot 2 = 1 + 2^n$   $\checkmark$

3. Let  $A = \{1, 2, 3\}$  and let  $R = \{[1, 3], [2, 2], [3, 1]\}$  be a relation on  $A$ . Find  $R \circ R$  and  $R \circ R \circ R$  and sketch a digraph for each of the relations  $R, R \circ R, R \circ R \circ R$

$$R \circ R = \{ [1, 1], [2, 2], [3, 3] \} = I_A, \text{ so } R \circ R \circ R = R$$



4. Define a relation  $R$  on  $\mathbf{R} \times \mathbf{R}$  by  $[x, y]R[r, s] \Leftrightarrow x + y = r + s$ . Prove that  $R$  is an equivalence relation. On the same set of axes sketch the equivalence class of  $[1, 2]$  and the equivalence class of  $[1, 3]$

**Reflexive:** if  $[x, y] = [r, s]$ , then  $x = r$  and  $y = s$ , so  $x + y = r + s$   $\checkmark$

**Symmetric:** by inspection  $\checkmark$


**Transitive:** if  $x + y = r + s$  and  $r + s = u + v$ , then  $x + y = u + v$   $\checkmark$

$$[x, y] \in [[1, 2]] \Leftrightarrow [x, y] R [1, 2] \Leftrightarrow x + y = 1 + 2 \Leftrightarrow y = -x + 3$$

$$\text{Thus, } [[1, 2]] = \{ [x, y] \in \mathbb{R} \times \mathbb{R} : y = -x + 3 \}$$

$$[x, y] \in [[1, 3]] \Leftrightarrow [x, y] R [1, 3] \Leftrightarrow x + y = 1 + 3 \Leftrightarrow y = -x + 4$$

$$\text{Thus, } [[1, 3]] = \{ [x, y] \in \mathbb{R} \times \mathbb{R} : y = -x + 4 \}$$



5. Explain why the set of all even integers  $2\mathbb{Z}$  and the set of all odd integers  $1 + 2\mathbb{Z}$  form a partition of  $\mathbb{Z}$ . Describe the equivalence relation on  $\mathbb{Z}$  whose quotient set is the above partition  $\{2\mathbb{Z}, 1 + 2\mathbb{Z}\}$

Cover:  $\mathbb{Z} \subseteq 2\mathbb{Z} \cup (1 + 2\mathbb{Z})$

If  $n \in \mathbb{Z}$ , then  $n$  is even or  $n$  is odd

Proof: By the Division Algorithm

if  $n \in \mathbb{Z}$ , then  $\exists! q, r \quad n = 2q + r \wedge 0 \leq r < 2 \quad (r = 0 \text{ or } 1)$

If  $r = 0$ , then  $n = 2q \in 2\mathbb{Z}$ . If  $r = 1$ , then  $n = 2q + 1 \in 1 + 2\mathbb{Z}$

so  $n \in 2\mathbb{Z} \cup (1 + 2\mathbb{Z}) \quad \ddot{\smile}$

Pairwise disjoint:  $2\mathbb{Z} \cap (1 + 2\mathbb{Z}) = \emptyset$

If  $n \in \mathbb{Z}$ , then  $n$  cannot be both even and odd

Proof: If  $2\mathbb{Z} \cap (1 + 2\mathbb{Z}) \neq \emptyset$ , then  $\exists n \in 2\mathbb{Z} \cap (1 + 2\mathbb{Z})$

Then  $n \in 2\mathbb{Z}$ , so  $\exists q_1 \in \mathbb{Z} \quad n = 2q_1$  and  $n \in 1 + 2\mathbb{Z}$ ,

so  $\exists q_2 \in \mathbb{Z} \quad n = 1 + 2q_2$ , so  $2q_1 = 1 + 2q_2$ , so  $1 = 2(q_1 - q_2) \quad \ddot{\smile}$

Suppose  $R$  is an equivalence relation on  $\mathbb{Z}$

such that  $\mathbb{Z}/R = \{2\mathbb{Z}, 1 + 2\mathbb{Z}\}$ .

(in other words the equivalence classes of  $R$  are  $2\mathbb{Z}$  and  $1 + 2\mathbb{Z}$ )

If  $m, n \in \mathbb{Z}$ , then

$$m R n \Leftrightarrow (m \in 2\mathbb{Z} \wedge n \in 2\mathbb{Z}) \vee (m \in 1 + 2\mathbb{Z} \wedge n \in 1 + 2\mathbb{Z})$$

$$\Leftrightarrow m \text{ and } n \text{ are even or } m \text{ and } n \text{ are odd}$$

$$\Leftrightarrow m \text{ and } n \text{ have the same parity} \quad \ddot{\smile}$$